

## An integrative model of shared decision making in medical encounters

Gregory Makoul\*, Marla L. Clayman

Program in Communication and Medicine, Division of General Internal Medicine,  
Robert H. Lurie Comprehensive Cancer Center, Northwestern University Feinberg School of Medicine,  
676 North St. Clair, Suite 200, Chicago, IL 60611, USA

Received 16 April 2005; received in revised form 6 June 2005; accepted 8 June 2005

### Abstract

**Objective:** Given the fluidity with which the term shared decision making (SDM) is used in teaching, assessment and research, we conducted a focused and systematic review of articles that specifically address SDM to determine the range of conceptual definitions.

**Methods:** In April 2005, we ran a Pubmed (Medline) search to identify articles published through 31 December 2003 with the words *shared decision making* in the title or abstract. The search yielded 681 citations, 342 of which were about SDM in the context of physician–patient encounters and published in English. We read and reviewed the full text of all 342 articles, and got any non-redundant references to SDM, which yielded an additional 76 articles.

**Results:** Of the 418 articles examined, 161 (38.5%) had a conceptual definition of SDM. We identified 31 separate concepts used to explicate SDM, but only “patient values/preferences” (67.1%) and “options” (50.9%) appeared in more than half the 161 definitions. Relatively few articles explicitly recognized and integrated previous work.

**Conclusion:** Our review reveals that there is no shared definition of SDM. We propose a definition that integrates the extant literature base and outlines essential elements that must be present for patients and providers to engage in the process of SDM.

**Practice implications:** The integrative definition of SDM is intended to provide a useful foundation for describing and operationalizing SDM in further research.

© 2005 Elsevier Ireland Ltd. All rights reserved.

**Keywords:** Shared decision making; Physician–patient relationship

### 1. Introduction

Decisions about tests, medications, procedures, referrals, or behaviors are an integral component of many medical encounters, and shared decision making (SDM) is frequently advocated in teaching and research about provider–patient interaction. However, the concept of SDM has been variably, and often loosely, defined. Some have acknowledged confusion surrounding the term [1–4], but recognition of the problem has not yet generated a model of SDM that integrates previous work. The lack of synthesis is problematic for several reasons. First, inconsistent conceptual definitions lead to inconsistent measurement of SDM

[1,4,5]. Second, the lack of a core definition of SDM complicates efforts to identify the relationships between SDM and outcome measures. Third, variable instantiations of SDM definitions make comparisons across studies difficult, if not impossible.

In terms of models of the provider–patient relationship, SDM is often positioned as a “middle ground” between paternalism (i.e., physicians make the decisions) and informed choice (i.e., patients make the decisions) [4,6–8]. In that context, there is considerable overlap between SDM and constructs with similar connotations, such as informed decision making [9], concordance [10,11], evidence-based patient choice [12,13], enhanced autonomy [14], and mutual participation [14]. There is a duality to the way SDM has been positioned within the proliferation of definitions. For instance, it has been described as both a

\* Corresponding author.

E-mail address: makoul@northwestern.edu (G. Makoul).

component of patient-centered care [15,16] and an extension of patient-centered medicine [17,18]. It has also been construed as the appropriate process for informed consent on one hand [19], and clearly distinguished from informed consent on the other [20–26].

Similarly, as noted by Charles et al. [1], models of SDM vary in the way they position the roles and responsibilities of each party. For example, Towle and Godolphin [27] suggested competencies for both physicians and patients, whereas others have placed more responsibility on the physician to elicit or respond to patients' views [28]. There has also been increasing attention to patients' preferred role in decision-making, with some asserting that for SDM to occur, patients must share equally in the decision-making process [4,7], while others contend that patients' role preferences be discussed and accepted [28].

### 1.1. Purpose of study

Given the fluidity with which the term *shared decision making* is used, we conducted a focused and systematic review of articles that specifically address SDM to determine the range of conceptual definitions therein. We sought to identify the most frequently invoked elements, qualities, and citations used to define SDM, with the goal of integrating the extant literature base to offer a conceptually sound and clinically relevant model of SDM.

## 2. Methods

In April 2005, we conducted a Pubmed (Medline) search to identify articles published through 31 December 2003 with the words *shared decision making* in the *title* or *abstract*.

### 2.1. Primary search strategy

The plan and procedure of our primary search strategy were designed to capture articles in the medical literature that clearly focus on shared decision making. We reasoned that if the words *shared decision making* appeared in the title or abstract, then SDM was likely to be a key topic or theme of the paper. While we were aware that the search strategy might miss some articles that addressed SDM in the text only [9,29–36], our intention was to examine articles that clearly purport to be about shared decision making. In addition, we were aware that some articles might not appear in Medline even if they include the words *shared decision making* in the title or abstract (e.g., Robert Kaplan's 1997 Presidential Address to the Society of Behavioral Medicine [37]) but maintained the straightforward search strategy in an effort to facilitate reproducibility. Similarly, we focused on articles published in English because their accessibility to the broad scientific community enhances reproducibility. This search yielded 681 articles that included the words *shared decision*




Fig. 1. Search strategy and results.

*making* in the title or abstract through 31 December 2003. At least two members of the study team reviewed the abstracts. As illustrated in Fig. 1, the next step in the review process was to drop any article that was not in English ( $n = 49$ ) or in which the words *shared decision making* were not in the context of the provider–patient relationship ( $n = 290$ ). We obtained the full text of all 342 articles that met the criteria for retention, including any articles that did not have an abstract (e.g., editorials).

### 2.2. Secondary search strategy

We read and reviewed the full text of all 342 articles, and got any non-redundant references to SDM whether the primary article included references within a conceptual definition of SDM or simply cited other articles after mentioning SDM. Consistent with our primary search strategy's focus on articles published in medically oriented journals, we did not obtain any referenced books, book chapters, or unpublished information such as course materials or technical reports. We conducted one iteration of this process, which yielded an additional 76 articles for subsequent coding (see Fig. 1).

### 2.3. Coding of conceptual definitions

Fig. 1 shows that our primary and secondary search strategies generated a total of 418 articles, 257 (61.5%) of which mentioned SDM without defining the term. Our coding of elements, qualities, and citations focuses on the remaining 161 articles that included a conceptual definition of SDM.

### 2.3.1. Elements and qualities of SDM

Each article was read by at least two members of the research team and sections that mentioned SDM were marked. We divided any definitions of SDM into units, which were generally delineated by conjunctions or punctuation. Then, based on a preliminary review of the unitized definitions, we developed coding rules and a coding sheet that included a list of mutually exclusive and exhaustive concepts (e.g., information exchange; patient preferences) plus an “other” category. Some of these coding categories represent elements of SDM (i.e., specific observable behaviors) while others describe qualities of SDM (i.e., relatively general characteristics). Conceptual definitions were coded by both authors, and the few discrepancies were resolved by reviewing together the articles in question. The specific content of any units coded as “other” was listed on the coding sheet; we created new coding categories for redundant “others”.

### 2.3.2. Citations about SDM

We also recorded all citations regarding SDM within the conceptual definitions to track how frequently


previously published articles and models were referenced. Our goal was to determine the extent to which conceptual definitions recognized, and were informed by, previous work.

## 3. Results

Of the 418 articles examined, 161 (38.5%) had a conceptual definition of SDM; the primary search strategy yielded 144 (42.1% of 342) articles with conceptual definitions [1–7,10–18,21,23–28,38–158], and the secondary search strategy yielded 17 (22.4% of 76) more [8,19,22,159–172]. Overall, 47.2% of the articles were classified as reviews/essays, 38.5% as empirical research reports, and 14.3% as editorials/commentaries.

### 3.1. Trajectory of the literature on SDM

The term “shared decision making” was first defined by the President’s Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral



All articles\* = results of the primary search strategy, which captured articles that were indexed in Pubmed (Medline) through 31 December 2003, included the words *shared decision making* in the title or abstract, and met the following two inclusion criteria: (1) in the context of patient-provider encounters; (2) published in English. N = 342.

Fig. 2. Growth in Pubmed-indexed articles on SDM.

Research in its report *Making Health Care Decisions*, which focused on informed consent [19]. This report was published in 1982, and cast SDM as a process based on mutual respect and partnership:

*It will usually consist of discussions between professional and patient that bring the knowledge, concerns, and perspective of each to the process of seeking agreement on a course of treatment. Simply put, this means that the physician or other health professional invites the patient to participate in a dialogue in which the professional seeks to help the patient understand the medical situation and available courses of action, and the patient conveys his or her concerns and wishes. This does not involve a mechanical recitation of abstruse medical information, but should include disclosures that give the patient an understanding of his or her condition and an appreciation of its consequences (p. 38). Shared decision making requires that a practitioner seek not only to understand each patient's needs and develop reasonable alternatives to meet those needs, but also to present the alternatives in a way that enables patients to choose one they prefer. To participate in this process, patients must engage in a dialogue with the practitioner and make their views on well-being clear (p. 44).*

This seminal report was found through our secondary search strategy (i.e., it was not indexed in Pubmed because it was issued as a governmental publication rather than a journal article). Fig. 2 focuses on articles that were indexed in Pubmed, and illustrates that relatively few included the concept of SDM until the late 1990s, when separate articles by Charles et al. [4], Coulter [52], and a report by Towle [173] appear to have triggered increased interest in, and publications about, SDM.

### 3.2. Concepts of SDM evident in the literature

Table 1 includes a list of elements and qualities that appeared in at least 10% of conceptual definitions of SDM. Of these 20 categories, only patient values/preferences (67.1%) and options (50.9%) appeared in more than half of the conceptual definitions. Eleven additional concepts are not listed in Table 1 because they appeared in fewer than 10% of definitions: involves at least two people (9.9%); recognize patient autonomy (8.7%); provide a comfortable environment (8.1%); arrange follow-up (6.8%); ascertain preferred format for information (6.2%); all parties have a legitimate interest in the decision (5.0%); uncertainty (5.0%); patient should understand enough to participate (3.7%); both physician and patient are experts (3.1%); division of labor between patient and physician (2.5%); a decision may require more than one visit (1.9%).

### 3.3. Prominent conceptual definitions in the literature

We also examined frequency of citations in order to identify the most prominent models of SDM, and Table 2

Table 1  
Concepts used in definitions of SDM<sup>a</sup>

67.1%	Patient values/preferences
50.9%	Options
46.0%	Partnership
37.3%	Patient participation
36.6%	Patient education
35.4%	Benefits/risks (pros/cons)
31.7%	Deliberation/negotiation
30.4%	Doctor knowledge/recommendations
29.2%	Mutual agreement
26.7%	Process/stages
23.6%	Middle ground
23.0%	Information exchange
18.0%	Make or explicitly defer decision
16.8%	Present evidence
13.0%	Define/explain problem
13.0%	Define roles (desire for involvement)
11.8%	Unbiased information
11.8%	Check/clarify understanding
11.2%	Flexibility/individualized approach
10.6%	Mutual respect

<sup>a</sup> Categories mentioned in at least 10% of articles with conceptual definitions are listed.

lists the authors of all models cited by more than 5% of articles that include conceptual definitions. Charles et al. [4] article was the most frequently cited, although it only appeared in 21.1% of the articles with conceptual definitions, illustrating that relatively few articles explicitly recognized and integrated previous work. Indeed, of the 161 articles with a conceptual definition of SDM, 56 (34.7%) did not cite any models. Within the most commonly cited articles, there has been a transition from identifying discrete characteristics of SDM [4] to focusing on process and outlining relatively sequential models [5,6,27,28,65,67,173].

## 4. Discussion and conclusion

Our review reveals that, overall, there is no shared definition of *shared decision making*. This is clearly the case within the set of articles that included a conceptual definition: We identified 31 separate concepts used to explicate SDM, only two of which appeared in more than half of the conceptual definitions. The lack of coherence looms even larger because 60% of articles that purport to focus on SDM failed to include any conceptual definition at

Table 2  
Most frequently cited definitions of SDM<sup>a</sup>

21.1%	Charles et al. [4]
9.9%	Towle and Godolphin [27]; Towle [173]
9.9%	Elwyn et al. [5,6,28,65–67]
8.7%	Charles et al. [7,8]
6.2%	President's Commission [19]
5.0%	Coulter and co-workers [52,54,162]

<sup>a</sup> Groups cited in at least 5% of articles with conceptual definitions are listed.

all. Equally troubling is the low frequency with which authors writing about SDM recognized and cited previous work in the area; fully one-third of articles with conceptual definitions failed to cite any other models. This proliferation of definitions can limit the productivity of research on SDM.

#### 4.1. Discussion

##### 4.1.1. Toward an integrative model

Thus, we propose an integrative model of SDM that builds upon the extant literature base. Our goal was to develop a definition that is conceptually sound, useful for future research, and applicable to clinical practice. We formulated this definition by taking the list of SDM elements (i.e., specific observable behaviors) and qualities (i.e., relatively general characteristics) listed in Table 1, and separating the elements into two groups: essential and ideal. More specifically, essential elements must be present for patients and providers to engage in the process of SDM; ideal elements may enhance the decision-making experience, but are not considered necessary for SDM to take place. We then identified which elements and qualities were positioned as central components of SDM within the prominent conceptual definitions listed in Table 2 (i.e., featured in focused paragraphs or lists). Finally, we added two elements: one that was cited in fewer than 10% of the

articles but evident in two of the relatively prominent models [5,6,27,28,65,67,173], and one that seems to be novel.

Table 3 displays this framework and lists the essential elements that comprise our integrative definition, illustrating patterns of emphasis in prominently cited models. In viewing the table, it is important to note that discussion of elements may be initiated by either physicians or patients. Accordingly, our choice of the term “elements” was a deliberate attempt to avoid placing sole responsibility on either party for displaying certain “competences” or “competencies” [81].

##### 4.1.2. Essential elements of SDM

In order for SDM to occur, patients and providers must first define and/or explain the problem that needs to be addressed. That discussion will likely lead to a presentation of options: Physicians should review options, if options exist, and patients should raise options of which they may be aware. Physicians and patients should discuss the pros and cons of options raised, particularly because they may have different perspectives on the relative importance of benefits, risks, and costs, including convenience and opportunity cost. These perspectives become evident through explication of patient values and preferences – including ideas, concerns, and outcome expectations – as well as physician knowledge and recommendations in the context of the decision at hand. Discussion of patients’ ability, or self-efficacy, to follow

Table 3  
Essential elements, ideal elements, and general qualities of SDM: emphasis in prominently cited models<sup>a</sup>

	President's Commission [19]	Charles et al. [4,7,8]	Coulter et al. [52,54,162]	Towle and Godolphin [27,173]	Elwyn et al. [5,6,28,65–67]
<b>Essential elements</b>					
Define/explain problem	X		X		X
Present options	X	X	X	X	X
Discuss pros/cons (benefits/risks/costs)		X	X	X	X
Patient values/preferences	X	X	X	X	X
Discuss patient ability/self-efficacy <sup>b</sup>					
Doctor knowledge/recommendations	X	X			
Check/clarify understanding		X			X
Make or explicitly defer decision		X	X	X	X
Arrange follow-up <sup>c</sup>				X	X
<b>Ideal elements</b>					
Unbiased information	X	X			X
Define roles (desire for involvement)		X	X	X	X
Present evidence		X	X	X	
Mutual agreement	X	X	X	X	X
<b>General qualities</b>					
Deliberation/negotiation	X	X		X	
Flexibility/individualized approach	X	X	X		X
Information exchange		X			
Involves at least two people		X		X	
Middle ground		X			
Mutual respect	X		X		
Partnership	X	X	X	X	
Patient education		X			X
Patient participation	X	X			X
Process/stages	X	X	X	X	X

<sup>a</sup> Unless otherwise indicated, this table includes categories from Table 1.

<sup>b</sup> This category was added by the authors.

<sup>c</sup> This category was cited in fewer than 10% of articles, so did not appear in Table 1.

through with a plan (e.g., test, medication, procedure, behavior change, referral) [174–178] is a critical – though often overlooked – component of assessing the viability of options. Throughout the process, both parties should periodically check understanding of facts and perspectives, providing further clarification as needed. The importance of checking and clarifying understanding has been reinforced by research on health literacy [179,180]. Of course, decisions are not always “made” when problems are first discussed; they may be explicitly deferred for a later time (e.g., pending discussion with members of the family and/or healthcare team) [4,7,18,93,154]. Thus, it is essential that physicians and patients arrange follow-up to track the outcome of decisions that have been made or reach resolution on those that have not.

There is considerable and intentional overlap between this integrative definition and previously published work. At least one of these elements is discussed in nearly all of the articles with conceptual definitions of SDM [1–8,10–19,21,23,24,26–28,38–49,51–56,58,59,61,62,64–69,72,74,76–78,81–86,88,90–97,101,102,105–108,110,113–117,120,121,123,124,126–131,133–135,139,141,142,144–146,148–150,152–160,162–165,171,172]. Taken together, these essential elements are rooted in the transactional model of communication, which holds that messages are filtered through different frames-of-reference, or life-spaces [181], and that people involved in an interaction simultaneously influence one another.

**4.1.2.1. Self-efficacy.** A brief overview of self-efficacy is warranted, given our suggestion that discussing patients’ ability to follow through with a plan be considered an essential element of SDM. The term self-efficacy refers to an individual’s own perceived ability to perform a specified behavior or set of behaviors. This is a construct central to Social Cognitive Theory, a variant of social learning theory that focuses on the continuous, reciprocal interaction between cognitions, behavior, and environment [174,175]. According to Social Cognitive Theory, behaviors are determined to a large extent by the outcome and efficacy expectations related to enacting them. Outcome expectations refer to an individual’s perceptions about whether behaviors will lead to certain outcomes (e.g., “taking this medicine every day will decrease my cholesterol”, “eating less salt will help reduce my blood pressure”). Efficacy expectations, often termed self-efficacy, refer to the individual’s beliefs about whether he or she can successfully enact the behavior in question (e.g., “I can take this medicine every day”, “I can eat less salt”). Individuals with greater perceived confidence with regard to a particular task, skill, or action may be more likely to engage in the behavior. Self-efficacy has been investigated in a variety of contexts and settings; it predicts the likelihood of initiating communication [182], adjusting to illness and treatment [183–187], adhering to medication treatment plans [188–190], and engaging in recommended health behaviors [191–

193]. In many ways, the rationale for incorporating a patient’s efficacy expectations parallels the argument for discussing patient preferences and values: both provide important perspective regarding acceptability of the options at hand.

#### 4.1.3. Ideal elements

Our integrative model of SDM is restricted to the essential elements because it is intended to encompass different clinical contexts, types of decisions and levels of involvement [6]. Some elements are considered “ideal” because they may enhance the process of SDM but are more applicable to some encounters than others. In other words, they are not always relevant or necessary for SDM to take place [9].

For instance, if it is indeed possible to deliver unbiased information, it may only be appropriate if the patient does not want the physician’s opinion [7,8,43,54]. Defining roles is another element considered ideal, particularly in light of Elwyn et al.’s argument that ascertaining role preference may be an implicit process [6]. Many authors have suggested that presentation of evidence is a key component of SDM [7,11,12,15,21,27,38,51,52,56,104,107,110,127,128,130,1–41,142,144,146,149,157], but we feel this must be considered an ideal element because adequate evidence is available for only a limited number of clinical decisions. Finally, while it has been suggested that a mutually agreed upon course of action is the appropriate result of SDM [27,52], a difference of opinion between physician and patient may still exist at the end of the SDM process [7,14,21]. We recognize that mutual agreement is highlighted in each of the prominent models, but believe it is properly positioned as an ideal rather than a necessity.

#### 4.1.4. General qualities

Extant conceptual definitions invoke several relatively general characteristics to describe SDM: deliberation/negotiation; flexibility/individualized approach; information exchange; involves at least two people; middle ground; mutual respect; partnership; patient education; patient participation; process/stages. While extremely useful in terms of providing an overall sense of SDM, these qualities do not highlight specific behaviors that readily lend themselves to research or practice [81]. However, they do raise issues that can help frame future research. For instance, the quality of “flexibility/individualized approach” suggests not only flexibility with respect to individual patients, but flexibility over time. In contexts such as cancer care and chronic disease management, physicians and patients may revisit decisions as circumstances change. Attention to this reality of everyday clinical practice sets the stage for longitudinal research on SDM.

#### 4.1.5. Can SDM be shared equally?

It is important to recognize that, while nearly half of the conceptual definitions invoked the concept of “partner-

ship", it is unlikely that decision-making will be shared equally in SDM. It may be helpful to envision the degree of sharing as a continuum with physicians leading the discussion and making decisions at one end, patients leading the discussion and making decisions at the other, and truly shared discussion and decision-making in the middle [194]. The nature of SDM will be qualitatively different as encounters depart from the midpoint: As illustrated in Fig. 3, whichever party leads the discussion, the degree of sharing increases as input from the other party increases (i.e., whether the other's acknowledgement, agreement, or opinion regarding the decision is either sought or offered). The balance of medical knowledge and social power in the provider–patient relationship is nearly always tipped toward physicians, and physicians often take a leadership role with respect to decisions in medical encounters [19]. Engaging in SDM does not require that physicians relinquish decision-making authority [52]. **Indeed, we contend that SDM can occur even if patients ask physicians to take decision-making responsibility, provided that the essential elements are present.**

#### 4.1.6. SDM and outcomes

Our study focuses on defining the concept of SDM, which is the prerequisite to addressing the question of whether SDM has positive outcomes. The answer to this question (i.e., empirical evidence) will likely determine how strongly the concept of SDM will be advocated and adopted in the years to come. Work by Greenfield and co-worker's research team links patient participation in medical care to improved health outcomes for patients with chronic illness [195–197]. In terms of measuring tangible and important outcomes, these studies offer a gold standard for research on patient participation. But they do not necessarily provide evidence about SDM because patient participation and SDM are not isomorphic. That is, patient participation in these studies stems from coaching patients to ask questions, negotiate

- doctor alone
- doctor led & patient acknowledgement (sought or offered)
- doctor led & patient agreement (sought or offered)
- doctor led & patient views/opinion (sought or offered)
- shared equally
- patient led & doctor views/opinion (sought or offered)
- patient led & doctor agreement (sought or offered)
- patient led & doctor acknowledgement (sought or offered)
- patient alone

Fig. 3. Degree of sharing.

medical decisions, and overcome barriers such as embarrassment and anxiety in the service of taking a more active role in their care [197]. This is a complex intervention, the components of which are difficult to disentangle. Future studies should build upon this base, focusing clearly on the extent to which implementing SDM improves health outcomes.

#### 4.2. Conclusion

A hallmark of shared decision making, that patients and providers have different – but equally valuable – perspectives and roles in the medical encounter was encapsulated in the title of a book published 20 years ago, *Meetings between Experts* [198]. This deceptively simple characterization provides a line of continuity throughout much of the literature on SDM, from the early conceptual definition in the President's Commission report through work published much more recently [14,17,19,116]. That said, the burgeoning interest in SDM requires a more clearly articulated and widely shared definition of the concept. Thus, we offer a definition that integrates the extant literature base while differentiating between essential elements (i.e., must be present for patients and providers to engage in the process of SDM) and ideal elements (i.e., may enhance the decision-making experience, but are not considered necessary for SDM to take place). There is considerable and intentional overlap between this integrative definition and previously published work.

#### 4.3. Practice implications

While SDM has received considerable attention in the context of difficult decisions (e.g., cancer treatment), it is equally important to study communication and decision-making in relatively mundane contexts such as primary care [9,18,19,27,159,199]. The integrative definition is intended to provide a useful foundation for describing and operationalizing SDM for further research on the meetings between experts in both pivotal circumstances and everyday clinical practice.

#### Acknowledgements

We are grateful to Amanda Zick (Program in Communication and Medicine, Northwestern University Feinberg School of Medicine, Chicago) and Rachel Malis (Department of Communication Studies, Northwestern University, Evanston) for their invaluable help with reviewing abstracts and articles. Drs. Cathy Charles and Aviram Gafni (McMaster University, Ontario), Angela Coulter (Picker Institute Europe, Oxford), Glyn Elwyn and Adrian Edwards (Cardiff University, Cardiff), and Angela Towle and William Godolphin (University of British Columbia, Vancouver) all graciously agreed to meet with us during the 2005

International Shared Decision Making Conference to discuss the integrative definition, and provided helpful input.

## References

- [1] Charles CA, Whelan T, Gafni A, Willan A, Farrell S. Shared treatment decision making: what does it mean to physicians? *J Clin Oncol* 2003;21:932–6.
- [2] Jansen LA. Deliberative decision making and the treatment of pain. *J Palliat Med* 2001;4:23–30.
- [3] Thornton H, Edwards A, Elwyn G. Evolving the multiple roles of 'patients' in health-care research: reflections after involvement in a trial of shared decision-making. *Health Expect* 2003;6:189–97.
- [4] Charles C, Gafni A, Whelan T. Shared decision-making in the medical encounter: what does it mean? (or it takes at least two to tango). *Soc Sci Med* 1997;44:681–92.
- [5] Elwyn G, Edwards A, Mowle S, Wensing M, Wilkinson C, Kinnersley P, Grol R. Measuring the involvement of patients in shared decision-making: a systematic review of instruments. *Patient Educ Couns* 2001;43:5–22.
- [6] Elwyn G, Edwards A, Gwyn R, Grol R. Towards a feasible model for shared decision making: focus group study with general practice registrars. *BMJ* 1999;319:753–6.
- [7] Charles C, Gafni A, Whelan T. Decision-making in the physician-patient encounter: revisiting the shared treatment decision-making model. *Soc Sci Med* 1999;49:651–61.
- [8] Charles C, Whelan T, Gafni A. What do we mean by partnership in making decisions about treatment? *BMJ* 1999;319:780–2.
- [9] Braddock III CH, Edwards KA, Hasenberg NM, Laidley TL, Levinson W. Informed decision making in outpatient practice: time to get back to basics. *JAMA* 1999;282:2313–20.
- [10] Taylor SG, Pickens JM, Geden EA. Interactional styles of nurse practitioners and physicians regarding patient decision making. *Nurs Res* 1989;38:50–5.
- [11] Jordan JL, Ellis SJ, Chambers R. Defining shared decision making and concordance: are they one and the same? *Postgrad Med J* 2002;78:383–4.
- [12] Ruland CM, Bakken S. Developing, implementing, and evaluating decision support systems for shared decision making in patient care: a conceptual model and case illustration. *J Biomed Inform* 2002;35:313–21.
- [13] Ford S, Schofield T, Hope T. What are the ingredients for a successful evidence-based patient choice consultation? A qualitative study. *Soc Sci Med* 2003;56:589–602.
- [14] Stubblefield C, Mutha S. Provider-patient roles in chronic disease management. *J Allied Health* 2002;31:87–92.
- [15] Godolphin W. The role of risk communication in shared decision making. *BMJ* 2003;327:692–3.
- [16] Teutsch C. Patient–doctor communication. *Med Clin North Am* 2003;87:1115–45.
- [17] Butler CC, Kinnersley P, Prout H, Rollnick S, Edwards A, Elwyn G. Antibiotics and shared decision-making in primary care. *J Antimicrob Chemother* 2001;48:435–40.
- [18] Stevenson FA, Barry CA, Britten N, Barber N, Bradley CP. Doctor-patient communication about drugs: the evidence for shared decision making. *Soc Sci Med* 2000;50:829–40.
- [19] President's Commission for the Study of Ethical Problems in Medicine and Biomedical and Behavioral Research. *Making Health Care Decisions. The Ethical and Legal Implications of Informed Consent in the Patient–Practitioner Relationship*. Washington, 1982.
- [20] Whitney SN, McGuire AL, McCollough LB. A typology of shared decision making, informed consent, and simple consent. *Ann Intern Med* 2004;140:54–9.
- [21] Hamann J, Leucht S, Kissling W. Shared decision making in psychiatry. *Acta Psychiatr Scand* 2003;107:403–9.
- [22] Smith DH, Pettegrew LS. Mutual persuasion as a model for doctor-patient communication. *Theor Med* 1986;7:127–46.
- [23] Frosch DL, Kaplan RM. Shared decision making in clinical medicine: past research and future directions. *Am J Prev Med* 1999;17:285–94.
- [24] Bartholome WG. A revolution in understanding: how ethics has transformed health care decision making. *QRB Qual Rev Bull* 1992;18:6–11.
- [25] Woolf SH. The logic and limits of shared decision making. *J Urol* 2001;166:244–5.
- [26] Deber RB. Shared decision making in the real world. *J Gen Intern Med* 1996;11:377–8.
- [27] Towle A, Godolphin W. Framework for teaching and learning informed shared decision making. *BMJ* 1999;319:766–71.
- [28] Elwyn G, Edwards A, Kinnersley P, Grol R. Shared decision making and the concept of equipoise: the competences of involving patients in healthcare choices. *Br J Gen Pract* 2000;50:892–9.
- [29] Cegala DJ, McClure L, Marinelli TM, Post DM. The effects of communication skills training on patients' participation during medical interviews. *Patient Educ Couns* 2000;41:209–22.
- [30] Donovan J. Patient decision making: the missing ingredient in compliance research. *Int J Technol Assess Health Care* 1995;11:443–55.
- [31] Edwards A, Elwyn G. The potential benefits of decision aids in clinical medicine. *JAMA* 1999;282:779–80.
- [32] Edwards A, Elwyn G. Understanding risk and lessons for clinical risk communication about treatment preferences. *Qual Health Care* 2001;10(Suppl. 1):i9–i13.
- [33] Holmes-Rovner M, Kroll J, Rovner DR, Schmitt N, Rothert M, Padonu G, Talarczyk G. Patient decision support intervention: increased consistency with decision analytic models. *Med Care* 1999;37:270–84.
- [34] Kaplan SH, Greenfield S, Gandek B, Rogers WH, Ware Jr JE. Characteristics of physicians with participatory decision-making styles. *Ann Intern Med* 1996;124:497–504.
- [35] Stewart MA. Effective physician–patient communication and health outcomes: a review. *CMAJ* 1995;152:1423–33.
- [36] Stiggelbout AM, Kiebert GM. A role for the sick role. Patient preferences regarding information and participation in clinical decision-making. *CMAJ* 1997;157:383–9.
- [37] Kaplan RM. Shared medical decision-making: a new paradigm for behavioral medicine—1997 presidential address. *Ann Behav Med* 1999;21:3–11.
- [38] Bauchner H. Shared decision making in pediatrics. *Arch Dis Child* 2001;84:246.
- [39] Bennett CL, Buchner DA, Ullman M. Approaches to prostate cancer by managed care organizations. *Urology* 1997;50:79–86.
- [40] Benowitz S. Shared decision-making still evolving in cancer screening. *J Natl Cancer Inst* 2001;93:672–3.
- [41] Bernat JL. Informed consent. *Muscle Nerve* 2001;24:614–21.
- [42] Brady TJ. The patient's role in rheumatology care. *Curr Opin Rheumatol* 1998;10:146–51.
- [43] Brock DW. The ideal of shared decision making between physicians and patients. *Kennedy Inst Ethic J* 1991;28–47.
- [44] Brown RF, Butow PN, Henman M, Dunn SM, Boyle F, Tattersall MH. Responding to the active and passive patient: flexibility is the key. *Health Expect* 2002;5:236–45.
- [45] Burke W, Beeker C, Kraft JM, Pinsky L. Engaging women's interest in colorectal cancer screening: a public health strategy. *J Womens Health Gend Based Med* 2000;9:363–71.
- [46] Burke W, Acheson L, Botkin J, Bridges K, Davis A, Evans J, Frias J, Hanson J, Kahn N, Kahn R, Lanier D, Pinsky LE, Press N, Lloyd-Puryear MA, Rich E, Stevens N, Thomson E, Wartman S, Wilson M. Genetics in primary care: a USA faculty development initiative. *Community Genet* 2002;5:138–46.
- [47] Carrere MO, Moumjid-Ferdjaoui N, Charavel M, Bremond A. Eliciting patients' preferences for adjuvant chemotherapy in breast

cancer: development and validation of a bedside decision-making instrument in a French Regional Cancer Centre. *Health Expect* 2000;3:97–113.

[48] Charavel M, Bremond A, Moumjid-Ferdjaoui N, Mignotte H, Carrere MO. Shared decision-making in question. *Psychooncology* 2001;10:93–102.

[49] Chin JJ. Doctor–patient relationship: from medical paternalism to enhanced autonomy. *Singapore Med J* 2002;43:152–5.

[50] Cohen H, Britten N. Who decides about prostate cancer treatment? A qualitative study. *Fam Pract* 2003;20:724–9.

[51] Coulter A. Assembling the evidence: patient-focused outcomes research. *Health Libr Rev* 1994;11:263–8.

[52] Coulter A. Partnerships with patients: the pros and cons of shared clinical decision-making. *J Health Serv Res Policy* 1997;2:112–21.

[53] Coulter A. Whatever happened to shared decision-making? *Health Expect* 2002;5:185–6.

[54] Coulter A. Patient information and shared decision-making in cancer care. *Br J Cancer* 2003;89(Suppl. 1):S15–6.

[55] Daley J, Forrow L. Ethical issues. *Prim Care* 1992;19:203–16.

[56] Davey P, Pagliari C, Hayes A. The patient's role in the spread and control of bacterial resistance to antibiotics. *Clin Microbiol Infect* 2002;8(Suppl. 2):43–68.

[57] Deber RB. Physicians in health care management. 8. The patient–physician partnership: decision making, problem solving and the desire to participate. *CMAJ* 1994;151:423–7.

[58] Edwards A, Evans R, Elwyn G. Manufactured but not imported: new directions for research in shared decision making support and skills. *Patient Educ Couns* 2003;50:33–8.

[59] Edwards A, Elwyn G, Hood K, Robling M, Atwell C, Holmes-Rovner M, Kinnersley P, Houston H, Russell I. The development of COMRADE—a patient-based outcome measure to evaluate the effectiveness of risk communication and treatment decision making in consultations. *Patient Educ Couns* 2003;50:311–22.

[60] Edwards A, Elwyn G, Smith C, Williams S, Thornton H. Consumers' views of quality in the consultation and their relevance to 'shared decision-making' approaches. *Health Expect* 2001;4:151–61.

[61] Elit L, Charles C, Gold I, Gafni A, Farrell S, Tedford S, Dal Bello D, Whelan T. Women's perceptions about treatment decision making for ovarian cancer. *Gynecol Oncol* 2003;88:89–95.

[62] Elwyn G, Edwards A, Britten N. "Doing prescribing": how might clinicians work differently for better, safer care. *Qual Saf Health Care* 2003;12(Suppl. 1):i33–6.

[63] Elwyn G, Edwards A, Eccles M, Rovner D. Decision analysis in patient care. *Lancet* 2001;358:571–4.

[64] Elwyn G, Edwards A, Kinnersley P. Shared decision-making in primary care: the neglected second half of the consultation. *Br J Gen Pract* 1999;49:477–82.

[65] Elwyn G, Edwards A, Wensing M, Hibbs R, Wilkinson C, Grol R. Shared decision making observed in clinical practice: visual displays of communication sequence and patterns. *J Eval Clin Pract* 2001;7: 211–21.

[66] Elwyn G, Edwards A, Wensing M, Hood K, Atwell C, Grol R. Shared decision making: developing the OPTION scale for measuring patient involvement. *Qual Saf Health Care* 2003;12:93–9.

[67] Elwyn G, Gray J, Clarke A. Shared decision making and non-directiveness in genetic counselling. *J Med Genet* 2000;37: 135–8.

[68] Elwyn G, Gwyn R, Edwards A, Grol R. Is 'shared decision-making' feasible in consultations for upper respiratory tract infections? Assessing the influence of antibiotic expectations using discourse analysis. *Health Expect* 1999;2:105–17.

[69] Emery J. Is informed choice in genetic testing a different breed of informed decision-making? A discussion paper. *Health Expect* 2001;4:81–6.

[70] Farrell MH, Murphy MA, Schneider CE. How underlying patient beliefs can affect physician–patient communication about prostate-specific antigen testing. *Eff Clin Pract* 2002;5:120–9.

[71] Fenton MV. Development of the scale of humanistic nursing behaviors. *Nurs Res* 1987;36:82–7.

[72] Fenton WS. Shared decision making: a model for the physician–patient relationship in the 21st century? *Acta Psychiatr Scand* 2003;107:401–2.

[73] Fillion E. How is medical decision-making shared? The case of haemophilia patients and doctors: the aftermath of the infected blood affair in France. *Health Expect* 2003;6:228–41.

[74] Frosch DL, Kaplan RM, Felitti V. The evaluation of two methods to facilitate shared decision making for men considering the prostate-specific antigen test. *J Gen Intern Med* 2001;16:391–8.

[75] Gafaranga J, Britten N. "Fire away": the opening sequence in general practice consultations. *Fam Pract* 2003;20:242–7.

[76] Gattellari M, Butow PN, Tattersall MH. Sharing decisions in cancer care. *Soc Sci Med* 2001;52:1865–78.

[77] Godolphin W, Towle A, McKendry R. Evaluation of the quality of patient information to support informed shared decision-making. *Health Expect* 2001;4:235–42.

[78] Goldring AB, Taylor SE, Kemeny ME, Anton PA. Impact of health beliefs, quality of life, and the physician–patient relationship on the treatment intentions of inflammatory bowel disease patients. *Health Psychol* 2002;21:219–28.

[79] Gomella LG, Albertsen PC, Benson MC, Forman JD, Soloway MS. The use of video-based patient education for shared decision-making in the treatment of prostate cancer. *Semin Urol Oncol* 2000;18: 182–7.

[80] Gramlich EP, Waitzfelder BE. Interactive video assists in clinical decision making. *Methods Inf Med* 1998;37:201–5.

[81] Greenhalgh T. Commentary: competencies for informed shared decision making. *BMJ* 1999;319:770.

[82] Guimond P, Bunn H, O'Connor AM, Jacobsen MJ, Tait VK, Drake ER, Graham ID, Stacey D, Elmslie T. Validation of a tool to assess health practitioners' decision support and communication skills. *Patient Educ Couns* 2003;50:235–45.

[83] Gurmankin AD, Baron J, Hershey JC, Ubel PA. The role of physicians' recommendations in medical treatment decisions. *Med Decis Making* 2002;22:262–71.

[84] Gwyn R, Elwyn G. When is a shared decision not (quite) a shared decision? Negotiating preferences in a general practice encounter. *Soc Sci Med* 1999;49:437–47.

[85] Gwyn R, Elwyn G, Edwards A, Mooney A. The problematic of decision-sharing: deconstructing 'cholesterol' in a clinical encounter. *Health Expect* 2003;6:242–54.

[86] Handley MR, Stuart ME. The use of prostate specific antigen for prostate cancer screening: a managed care perspective. *J Urol* 1994;152:1689–92.

[87] Healey JM. Futility, shared decision making, and the doctor–patient relationship. *Conn Med* 1992;56:166.

[88] Hibbard JH. Engaging health care consumers to improve the quality of care. *Med Care* 2003;41:61–70.

[89] Holmes-Rovner M, Valade D, Orlowski C, Draus C, Nabozny-Valerio B, Keiser S. Implementing shared decision-making in routine practice: barriers and opportunities. *Health Expect* 2000;3:182–91.

[90] Hudak PL, Clark JP, Hawker GA, Coyte PC, Mahomed NN, Kreder HJ, Wright JG. "You're perfect for the procedure! Why don't you want it?" Elderly arthritis patients' unwillingness to consider total joint arthroplasty surgery: a qualitative study. *Med Decis Making* 2002;22:272–8.

[91] Juniper EF. The impact of patient compliance on effective asthma management. *Curr Opin Pulm Med* 2003;9(Suppl. 1):S8–S10.

[92] Kaplan RM. The significance of quality of life in health care. *Qual Life Res* 2003;12:3–16.

[93] Karlawish JH. Shared decision making in critical care: a clinical reality and an ethical necessity. *Am J Crit Care* 1996;5:391–6.

[94] Kasper JF, Fowler Jr FJ. Responding to the challenge. A status report on shared decision-making programs. *HMO Pract* 1993;7:176–81.

[95] Kasper JF, Mulley Jr AG, Wennberg JE. Developing shared decision-making programs to improve the quality of health care. *QRB Qual Rev Bull* 1992;18:183–90.

[96] Keefe CW, Thompson ME, Noel MM. Medical students, clinical preventive services, and shared decision-making. *Acad Med* 2002;77:1160–1.

[97] Kerridge IH, Pearson SA, Rolfe IE, Lowe M. Decision making in CPR: attitudes of hospital patients and healthcare professionals. *Med J Aust* 1998;169:128–31.

[98] Kessler S. Psychological aspects of genetic counselling. XIV. Non-directiveness and counseling skills. *Genet Test* 2001;5:187–91.

[99] Kim SP, Knight SJ, Tomori C, Colella KM, Schoor RA, Shih L, Kuzel TM, Nadler RB, Bennett CL. Health literacy and shared decision making for prostate cancer patients with low socioeconomic status. *Cancer Invest* 2001;19:684–91.

[100] Koehle M, Loyd-Smith R, McKenzie D, Taunton J. Asthma and recreational SCUBA diving: a systematic review. *Sports Med* 2003;33:109–16.

[101] Leland J. Advance directives and establishing the goals of care. *Prim Care* 2001;28:349–63.

[102] Lelie A. Decision-making in nephrology: shared decision making? *Patient Educ Couns* 2000;39:81–9.

[103] Lenert LA, Cher DJ. Use of meta-analytic results to facilitate shared decision making. *J Am Med Inform Assoc* 1999;6:412–9.

[104] Liaw ST, Pearce C, Jackson B. Anticoagulant therapy: will computer use improve outcomes? *Aust Fam Physician* 2001;30:964–8.

[105] Loewy RS. Honouring the age-old commitment to “the patient’s good”: the promise-and peril-of hospice. *Wien Med Wochenschr* 2003;153:392–7.

[106] Lowance DC. Withholding and withdrawal of dialysis in the elderly. *Semin Dial* 2002;15:88–90.

[107] Lurie JD, Weinstein JN. Shared decision-making and the orthopaedic workforce. *Clin Orthop* 2001;68–75.

[108] Malm U, Ivarsson B, Allebeck P, Falloon IR. Integrated care in schizophrenia: a 2-year randomized controlled study of two community-based treatment programs. *Acta Psychiatr Scand* 2003;107:415–23.

[109] McAlister FA. Atrial fibrillation, shared decision making, and the prevention of stroke. *Stroke* 2002;33:238–42.

[110] McHugo GJ, Drake RE. Finding and evaluating the evidence: a critical step in evidence-based medicine. *Psychiatr Clin North Am* 2003;26:821–31.

[111] McKeown RE, Reininger BM, Martin M, Hoppmann RA. Shared decision making: views of first-year residents and clinic patients. *Acad Med* 2002;77:438–45.

[112] Miller PA, Forbes S, Boyle DK. End-of-life care in the intensive care unit: a challenge for nurses. *Am J Crit Care* 2001;10:230–7.

[113] Mohr C, Milgrom J, Griffiths M, Nomikoudis K. Breaking the bad news: dilemmas in shared decision-making in medical practice.. *Aust Psychol* 1999;34:45–8.

[114] Montgomery AA, Fahey T. How do patients’ treatment preferences compare with those of clinicians? *Qual Health Care* 2001;10(Suppl. 1):i39–43.

[115] Morgan MW, Deber RB, Llewellyn-Thomas HA, Gladstone P, Cusimano RJ, O’Rourke K, Tomlinson G, Detsky AS. Randomized, controlled trial of an interactive videodisc decision aid for patients with ischemic heart disease. *J Gen Intern Med* 2000;15:685–93.

[116] Moss AH. Shared decision-making in dialysis: the new RPA/ASN guideline on appropriate initiation and withdrawal of treatment. *Am J Kidney Dis* 2001;37:1081–91.

[117] Moumid N, Bremond A, Carrere MO. From information to shared decision-making in medicine. *Health Expect* 2003;6:187–8.

[118] Myers RE, Kunkel EJ. Preparatory education for informed decision-making in prostate cancer early detection and treatment. *Semin Urol Oncol* 2000;18:172–7.

[119] Parascandola M, Hawkins J, Danis M. Patient autonomy and the challenge of clinical uncertainty. *Kennedy Inst Ethic J* 2002;12:245–64.

[120] Piercy GB, Deber R, Trachtenberg J, Ramsey EW, Norman RW, Goldenberg SL, Nickle JC, Elhilali M, Perrault JP, Kraetschmer N, Sharpe N. Impact of a shared decision-making program on patients with benign prostatic hyperplasia. *Urology* 1999;53:913–20.

[121] Pignone M, Bucholtz D, Harris R. Patient preferences for colon cancer screening. *J Gen Intern Med* 1999;14:432–7.

[122] Plumeri PA. Informed consent and the gastrointestinal endoscopist. *Gastrointest Endosc* 1985;31:218–21.

[123] Prout MN. Breast cancer risk reduction: what do we know and where should we go? *Medscape Womens Health* 2000;5:E4.

[124] Robinson A, Thomson R. Variability in patient preferences for participating in medical decision making: implication for the use of decision support tools. *Qual Health Care* 2001;10(Suppl. 1):i34–8.

[125] Rockefeller R. Informed shared decision making: is this the future of health care? *Health Forum J* 1999;42:54–6.

[126] Rothert ML, O’Connor AM. Health decisions and decision support for women. *Ann Rev Nurs Res* 2001;19:307–24.

[127] Rubin GL, Frommer MS, Vincent NC, Phillips PA, Leeder SR. Getting new evidence into medicine. *Med J Aust* 2000;172:180–3.

[128] Ruland CM. Handheld technology to improve patient care: evaluating a support system for preference-based care planning at the bedside. *J Am Med Inform Assoc* 2002;9:192–201.

[129] Ruland CM, Bakken S. Representing patient preference-related concepts for inclusion in electronic health records. *J Biomed Inform* 2001;34:415–22.

[130] Ruland CM, White T, Stevens M, Fanciullo G, Khilani SM. Effects of a computerized system to support shared decision making in symptom management of cancer patients: preliminary results. *J Am Med Inform Assoc* 2003;10:573–9.

[131] Rushton CH, Glover JJ. Involving parents in decisions to forego life-sustaining treatment for critically ill infants and children. *AACN Clin Issues Crit Care Nurs* 1990;1:206–14.

[132] Schofield T, Elwyn G, Edwards A, Visser A. Shared decision making. *Patient Educ Couns* 2003;50:229–30.

[133] Sculpher M, Gafni A, Watt I. Shared treatment decision making in a collectively funded health care system: possible conflicts and some potential solutions. *Soc Sci Med* 2002;54:1369–77.

[134] So A, Goldenberg L, Gleave ME. Prostate specific antigen: an updated review. *Can J Urol* 2003;10:2040–50.

[135] Stalmeier PF, Unic IJ, Verhoef LC, van Daal WA. Evaluation of a shared decision making program for women suspected to have a genetic predisposition to breast cancer: preliminary results. *Med Decis Making* 1999;19:230–41.

[136] Steinhart B. Patient autonomy: evolution of the doctor–patient relationship. *Haemophilia* 2002;8:441–6.

[137] Steven K. Shared decision-making. *Br J Gen Pract* 2001;51:61–2.

[138] Stevenson FA. General practitioners’ views on shared decision making: a qualitative analysis. *Patient Educ Couns* 2003;50:291–3.

[139] Stiggelbout AM, de Haes JC. Patient preference for cancer therapy: an overview of measurement approaches. *J Clin Oncol* 2001;19:220–30.

[140] Stutts A, Schloemann J. Life-sustaining support: ethical, cultural, and spiritual conflicts. Part I. Family support—a neonatal case study. *Neonat Netw* 2002;21:23–9.

[141] Sullivan KE, Hebert PC, Logan J, O’Connor AM, McNeely PD. What do physicians tell patients with end-stage COPD about intubation and mechanical ventilation? *Chest* 1996;109:258–64.

[142] Towle A. Shifting the culture of continuing medical education: what needs to happen and why is it so difficult? *J Contin Educ Health Prof* 2000;20:208–18.

[143] Vandevusse L. Decision making in analyses of women’s birth stories. *Birth* 1999;26:43–50.

[144] Volk RJ, Cass AR, Spann SJ. A randomized controlled trial of shared decision making for prostate cancer screening. *Arch Fam Med* 1999;8:333–40.

[145] Walker RM. Ethical issues in end-of-life care. *Cancer Control* 1999;6:162–7.

[146] Wei JT, Uzzo RG. Shared decision-making strategies for early prostate cancer. *Semin Urol Oncol* 2002;20:74–8.

[147] Wennberg JE, Fisher ES, Skinner JS. Geography and the debate over medicare reform. *Health Affair* 2002;96–114. Supp Web Exclusives.

[148] Weinstock MP. Shared decision-making makes medical paternalism obsolete. *So long, Marcus Welby*. *Volunt Leader* 1998;39: 11.

[149] Wensing M, Elwyn G, Edwards A, Vingerhoets E, Grol R. Deconstructing patient centred communication and uncovering shared decision making: an observational study. *BMC Med Inform Decis Mak* 2002;2:2.

[150] Whelan T, Levine MN, Gafni A, Sanders K, Willan A, Mirsky D, Schnider D, McCready D, Reid S, Kobylecky A, Reed K. Mastectomy or lumpectomy? Helping women make informed choices. *J Clin Oncol* 1999;17:1727–35.

[151] Whelan TJ, Levine MN, Gafni A, Lukka H, Mohide EA, Patel M, Streiner DL. Breast irradiation postlumpectomy: development and evaluation of a decision instrument. *J Clin Oncol* 1995;13:847–53.

[152] Whelan T, Gafni A, Charles C, Levine M. Lessons learned from the decision board: a unique and evolving decision aid. *Health Expect* 2000;3:69–76.

[153] Whitney SN. A new model of medical decisions: exploring the limits of shared decision making. *Med Decis Making* 2003;23:275–80.

[154] Wilkinson CR, Williams M. Strengthening patient–provider relationships. *Lippincott Case Manag* 2002;7:86–99.

[155] Wills CE, Holmes-Rovner M. Patient comprehension of information for shared treatment decision making: state of the art and future directions. *Patient Educ Couns* 2003;50:285–90.

[156] Wilt TJ, Partin MR. Prostate cancer intervention. Involving the patient in early detection and treatment. *Postgrad Med* 2003; 114:43–9.

[157] Woolf SH. Shared decision-making: the case for letting patients decide which choice is best. *J Fam Pract* 1997;45:205–8.

[158] Yen JC, Neville C, Fortin PR. Discordance between patients and their physicians in the assessment of lupus disease activity: relevance for clinical trials. *Lupus* 1999;8:660–70.

[159] Braddock III CH, Fihn SD, Levinson W, Jonsen AR, Pearlman RA. How doctors and patients discuss routine clinical decisions. Informed decision making in the outpatient setting. *J Gen Intern Med* 1997;12:339–45.

[160] Brock DW, Wartman SA. When competent patients make irrational choices. *N Engl J Med* 1990;322:1595–9.

[161] Brody DS, Miller SM, Lerman CE, Smith DG, Caputo GC. Patient perception of involvement in medical care: relationship to illness attitudes and outcomes. *J Gen Intern Med* 1989;4:506–11.

[162] Coulter A, Entwistle V, Gilbert D. Sharing decisions with patients: is the information good enough? *BMJ* 1999;318:318–22.

[163] Deber RB. Physicians in health care management. 7. The patient–physician partnership: changing roles and the desire for information. *CMAJ* 1994;151:171–6.

[164] Emanuel EJ, Emanuel LL. Four models of the physician–patient relationship. *JAMA* 1992;267:2221–6.

[165] Entwistle VJS, Watt I. Evaluating interventions to promote patient involvement in decision-making: by what criteria should effectiveness be judged? *J Health Serv Res Policy* 1998;3:100–7.

[166] Forrow L, Wartman SA, Brock DW. Science, ethics, and the making of clinical decisions. Implications for risk factor intervention. *JAMA* 1988;259:3161–7.

[167] Levine MN, Gafni A, Markham B, MacFarlane D. A bedside decision instrument to elicit a patient's preference concerning adjuvant chemotherapy for breast cancer. *Ann Intern Med* 1992;117:53–8.

[168] Molenaar S, Sprangers MA, Postma-Schuit FC, Rutgers EJ, Noorderlander J, Hendriks J, de Haes HC. Feasibility and effects of decision aids. *Med Decis Making* 2000;20:112–27.

[169] Rees AM. Communication in the physician–patient relationship. *Bull Med Libr Assoc* 1993;81:1–10.

[170] Sebba C, Brownman G, Gafni A, Norman G, Levine M, Assouline D, Fiere D. Design and validation of a bedside decision instrument to elicit a patient's preference concerning allogenic bone marrow transplantation in chronic myeloid leukemia. *Am J Hematol* 1995;48:221–7.

[171] Smith RA, von Eschenbach AC, Wender R, Levin B, Byers T, Rothenberger D, Brooks D, Creasman W, Cohen C, Runowicz C, Saslow D, Cokkinides V, Eyre H. American Cancer Society guidelines for the early detection of cancer: update of early detection guidelines for prostate, colorectal, and endometrial cancers. Also: update 2001—testing for early lung cancer detection. *CA Cancer J Clin* 2001;51:38–75.

[172] Wennberg JE. Outcomes research, cost containment, and fear of health care rationing. *N Eng J Med* 1990;323:1202–4.

[173] Towle A. Physician and patient communication skills: competencies for informed shared decision making. Vancouver, Canada: University of British Columbia, 1997.

[174] Bandura A. Self-efficacy: toward a unifying theory of behavioral change. *Psychol Rev* 1977;84:191–215.

[175] Bandura A. A social foundation of thought and action. A social cognitive theory. Englewood Cliffs, NJ: Prentice Hall, 1986.

[176] Bandura A. Reflections on self efficacy. *Adv Behav Res Ther* 1978;1:237–69.

[177] Bandura A. Self-efficacy mechanisms in human agency. *Am Psychol* 1982;37:122–47.

[178] Bandura A. Self-efficacy conception of anxiety. *Anxiety Res* 1988; 1.

[179] Gazmararian JA, Williams MV, Peel J, Baker DW. Health literacy and knowledge of chronic disease. *Patient Educ Couns* 2003;51:267–75.

[180] Wolf MS, Davis TC, Cross JT, Marin E, Green K, Bennett CL. Health literacy and patient knowledge in a Southern US HIV clinic. *Int J STD AIDS* 2004;15:747–52.

[181] Lewin K. A dynamic theory of personality. New York: McGraw-Hill, 1935.

[182] Makoul G, Roloff ME. The role of efficacy and outcome expectations in the decision to withhold relational complaints. *Commun Res* 1998;25:5–29.

[183] Bodenheimer T, Lorig K, Holman H, Grumbach K. Patient self-management of chronic disease in primary care. *JAMA* 2002;288: 2469–75.

[184] Lorig K, Chastain RL, Ung E, Shoor S, Holman HR. Development and evaluation of a scale to measure perceived self-efficacy in people with arthritis. *Arthritis Rheum* 1989;32:37–44.

[185] Lorig KR, Sobel DS, Stewart AL, Brown Jr BW, Bandura A, Ritter P, Gonzalez VM, Laurent DD, Holman HR. Evidence suggesting that a chronic disease self-management program can improve health status while reducing hospitalization: a randomized trial. *Med Care* 1999;37:5–14.

[186] Airlie J, Baker GA, Smith SJ, Young CA. Measuring the impact of multiple sclerosis on psychosocial functioning: the development of a new self-efficacy scale. *Clin Rehabil* 2001;15:259–65.

[187] Hewlett S, Cockshott Z, Kirwan J, Barrett J, Stamp J, Haslock I. Development and validation of a self-efficacy scale for use in British patients with rheumatoid arthritis (RASE). *Rheumatology (Oxford)* 2001;40:1221–30.

[188] Ogedegbe G, Mancuso CA, Allegrante JP, Charlson ME. Development and evaluation of a medication adherence self-efficacy scale in hypertensive African–American patients. *J Clin Epidemiol* 2003;56: 520–9.

[189] Scherer YK, Bruce S. Knowledge, attitudes, and self-efficacy and compliance with medical regimen, number of emergency department

visits, and hospitalizations in adults with asthma. *Heart Lung* 2001;30:250–7.

[190] Brus H, van de Laar M, Taal E, Rasker J, Wiegman O. Determinants of compliance with medication in patients with rheumatoid arthritis: the importance of self-efficacy expectations. *Patient Educ Couns* 1999;36:57–64.

[191] Marcus BH, Selby VC, Niaura RS, Rossi JS. Self-efficacy and the stages of exercise behavior change. *Res Q Exerc Sport* 1992;63:60–6.

[192] Forsyth AD, Carey MP. Measuring self-efficacy in the context of HIV risk reduction: research challenges and recommendations. *Health Psychol* 1998;17:559–68.

[193] Etter JF, Bergman MM, Humair JP, Perneger TV. Development and validation of a scale measuring self-efficacy of current and former smokers. *Addiction* 2000;95:901–13.

[194] Makoul G. Evidence based patient choice: theory vs. practice. *J Gen Intern Med* 2003;18S:211.

[195] Greenfield S, Kaplan S, Ware Jr JE. Expanding patient involvement in care. Effects on patient outcomes. *Ann Intern Med* 1985;102:520–8.

[196] Greenfield S, Kaplan SH, Ware Jr JE, Yano EM, Frank HJ. Patients' participation in medical care: effects on blood sugar control and quality of life in diabetes. *J Gen Intern Med* 1988;3:448–57.

[197] Kaplan SH, Greenfield S, Ware Jr JE. Assessing the effects of physician–patient interactions on the outcomes of chronic disease. *Med Care* 1989;27:S110–27.

[198] Tuckett D, Boulton M, Olson C, Williams A. Meetings between experts: an approach to sharing ideas in medical consultations.. New York: Tavistock Publications Ltd., 1985.

[199] Makoul G, Arntson P, Schofield T. Health promotion in primary care: physician–patient communication and decision making about prescription medications. *Soc Sci Med* 1995;41:1241–54.